MAR 19680106: NORTHWEST ALBERTA

Received date: Dec 31, 1968

Public release date: Jan 01, 1970

DISCLAIMER

By accessing and using the Alberta Energy website to download or otherwise obtain a scanned mineral assessment report, you ("User") agree to be bound by the following terms and conditions:

- a) Each scanned mineral assessment report that is downloaded or otherwise obtained from Alberta Energy is provided "AS IS", with no warranties or representations of any kind whatsoever from Her Majesty the Queen in Right of Alberta, as represented by the Minister of Energy ("Minister"), expressed or implied, including, but not limited to, no warranties or other representations from the Minister, regarding the content, accuracy, reliability, use or results from the use of or the integrity, completeness, quality or legibility of each such scanned mineral assessment report;
- b) To the fullest extent permitted by applicable laws, the Minister hereby expressly disclaims, and is released from, liability and responsibility for all warranties and conditions, expressed or implied, in relation to each scanned mineral assessment report shown or displayed on the Alberta Energy website including but not limited to warranties as to the satisfactory quality of or the fitness of the scanned mineral assessment report for a particular purpose and warranties as to the non-infringement or other non-violation of the proprietary rights held by any third party in respect of the scanned mineral assessment report:
- c) To the fullest extent permitted by applicable law, the Minister, and the Minister's employees and agents, exclude and disclaim liability to the User for losses and damages of whatsoever nature and howsoever arising including, without limitation, any direct, indirect, special, consequential, punitive or incidental damages, loss of use, loss of data, loss caused by a virus, loss of income or profit, claims of third parties, even if Alberta Energy have been advised of the possibility of such damages or losses, arising out of or in connection with the use of the Alberta Energy website, including the accessing or downloading of the scanned mineral assessment report and the use for any purpose of the scanned mineral assessment report so downloaded or retrieved.
- d) User agrees to indemnify and hold harmless the Minister, and the Minister's employees and agents against and from any and all third party claims, losses, liabilities, demands, actions or proceedings related to the downloading, distribution, transmissions, storage, redistribution, reproduction or exploitation of each scanned mineral assessment report obtained by the User from Alberta Energy.



ECONOMIC MINERALS

FILE REPORT No.

S-AF-126(1)

GENERAL GEOLOGY AND SULPHUR PROSPECTS

OF

SULPHUR PROSPECTING PERMIT NO. 126

FOR

JOSEPH ABRAHAM TANNOUS

BY

WILLIAM G. CROOK, P. GEOL.

INDEX

		1	110
LOCATION AND	ACCESS		1
STRATIGRAPHY			2
SOURCES OF SU	JLPHUR		7


LOCATION AND ACCESS


Sulphur Prospecting Permit No. 126 is located in Township 112, Range 9, West of the Fifth Meridian. This is in northwest Alberta, 335 miles north of Edmonton, 85 miles south of the Northwest Territories-Alberta border and 175 miles east of the British Columbia-Alberta border.


The Permit can be reached by car from Peace River via the MacKenzie Highway to High Level, then east on a secondary road to Fort Vermilion. From here several seismic roads and wellsite access roads lead east to the area of the Permit. These latter roads can only be used during the winter months when the ground is frozen.

A float-equipped aircraft can land on the lakes near the Permit.

Figure 1.







STRATIGRAPHY

The sedimentary section under Sulphur Permit No. 126 is about 5,000 feet thick, and the Devonian and Cretaceous systems are represented. There are four large unconformities in the section. They are, from older to younger:

- 1. Between the Pre-Cambrian Basement and the Middle Devonian.
- Between the Middle Devonian and the Upper Devonian.
- 3. Between the Upper Devonian and the Cretaceous.
- 4. Between the Cretaceous and the Tertiary.

The discussion of the stratigraphy will be based on the section as found at Crusade Dynamic Caribou 7-8-113-5-W.5 This well was drilled to the Red Beds formation and is located 18 miles northeast of the Permit.

DEVONIAN

The Devonian is about 2,760 feet thick and both Middle and Upper Devonian are present.

MIDDLE DEVONIAN

The Middle Devonian is 1,490 feet thick and may be assigned to the Elk Point Group.

ELK POINT GROUP

The Elk Point may be divided into five formations, viz - Red Beds, Chinchaga, Keg River, Muskeg and Watt Mountain.

RED BEDS FORMATION

The Red Beds are about 340 feet thick and consists of orange and red (rarely green and brown) anhydrite, marl and dolomite. Salt casts are common. The Granite Wash sandstone often occurs at the base of the Red Beds, and where present is a fine to coarse grained arkozic sandstone.

CHINCHAGA FORMATION

The Chinchaga is about 185 feet thick and is composed of brown, dense, anhydrite, with thin beds of brown microgranular and granular dolomite.

KEG RIVER FORMATION

The Keg River is 115 feet thick and consists of light brown to white, massive anhydrite, with thin beds of fine crystalline evaporitic dolomite.

MUSKEG FORMATION

The Muskeg is 810 feet thick and consists of light brown to white, massive anhydrite, with thin beds of fine crystalline evaporitic dolomite.

WATT MOUNTAIN FORMATION

The Watt Mountain is 40 feet thick and consists of green, black and brown waxy shale, with some anhydrite and dolomite.

UPPER DEVONIAN

The Upper Devonian is 1,280 feet thick and the following units are present, in ascending order - Slave Point and Shale Unit.

SLAVE POINT

The Slave Point is 148 feet thick and consists of an upper 62 feet of brown, granular to crystalline, fossiliferous limestone; and a lower 86 feet of white to brown massive anhydrite with some gray to buff, evaporitic dolomite (Fort Vermilion member).

SHALE UNIT

The Shale Unit is about 1,130 feet thick and consists of green and brown, calcareous, splintery shale. The shale is often silty and thin beds of light gray-brown limestone are

common near the base of the unit. The Shale Unit correlates with the Beaverhill Lake and the Hay River Shale.

CRETACEOUS

The Cretaceous section is about 2,300 feet thick at Crusade Dynamic Caribou 7-8 and both Lower and Upper Cretaceous units are present.

LOWER CRETACEOUS

The Lower Cretaceous is 1,230 feet thick and consists of the Spirit River and Peace River formations. The Spirit River is about 1,150 feet thick and consists of gray, sandy, chunky shale with thin beds and stringers of sandstone. The Peace River is about 80 feet thick and consists of the Cadotte sand which is 60 feet thick and the Harmon Shale which is about 20 feet thick.

UPPER CRETACEOUS

The Upper Cretaceous is about 1,000 feet thick and consists of the Shaftesbury Shale (700 feet thick) and the Dunvegan formation (300 feet thick) which consists of interbedded sandstone and shale.

TERTIARY

PLEISTOCENE

A layer of sand, gravel, clay and till of glacial origin lies on the surface of the Permit. It has a variable thickness but should not exceed 60 feet.

SOURCES OF SULPHUR

The sulphur bearing horizons under the Permit may be classed as follows:

CRETACEOUS

The writer has examined the Cretaceous shales in nearby outcrops and without exception the outcrops do contain small amounts of sulphur, plus greater amounts of selenite and ironstone concretions. However, these sulphur beds are so thin (maximum one-eighth of a inch thick) that they are of no commercial value. Usually the sulphur consists of yellow streaks on the shale.

DEVONIAN

Some minor traces of elemental sulphur has been noted in the Grosmont (D-2) in the Red Earth Oil Field 165 miles to the south, but no sulphur has been reported in well cuttings from tests near this permit.

SULPHUR WATER

The porous horizons within the Devonian often contain salty sulphur water. At the moment there is no commercial method of extracting the sulphur from the water but, undoubtedly such an extraction process will be developed in the future.

In conclusion it can be briefly stated that the prospects of finding elemental sulphur in commercial quantities under the Permit are very remote. Therefore, it is my recommendation that the Permit be dropped.

Respectfully submitted by:

WILLIAM G. CROOK, P. GEOL.

WGC/jp

SULPHUR PROSPECTING PERMIT No. 126

